

Vivendi[®] 200

MAPP 15017

Soluble Concentrate (SL) containing 200 g/l (18.2 % w/w) clopyralid.

VIVENDI 200 is a post-emergence foliar acting herbicide for use in many arable, forage and horticultural crops for the control of some annual broad-leaved weeds, but especially for the control of thistles.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

IMPORTANT INFORMATION

FOR USE AS AN AGRICULTURAL/HORTICULTURAL HERBICIDE

Crops	Maximum individual dose (litres product/hectare)	Maximum total dose (litres product/ha/crop)	Latest time of application
Winter and spring wheat, winter and spring barley and oats	0.35	0.35	before the third node is detectable
Oilseed rape	1.0	1.5	before flower buds are visible from above the crop canopy
Sugar beet, fodder beet, red beet and mangels	1.0	1.5	when the crops meet between the rows (BBCH 39)
Swede, turnip	1.0	1.5	when the crops meet between the rows (BBCH 39)
Permanent grassland, rotational grass	1.0	1.0 product/ha/year	7 days before grazing or cutting for hay and silage
Ornamental plant production	1.0	1.0 product/ha/year	-

Other specific restrictions Applications must not be made earlier than 1 March in the year of harvest. KEEP LIVESTOCK out of treated areas for at least 7 days and until foliage of any poisonous weeds such as ragwort have died and become unpalatable.
For use on ornamentals application must be only via directed sprays only to target weed foliage. Do not use between the end of August and beginning of March.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

SAFETY PRECAUTIONS

To avoid risks to man and the environment, comply with the instructions for use.

Operator protection

WHEN USING DO NOT EAT, DRINK OR SMOKE

DO NOT BREATHE SPRAY.

WASH CONCENTRATE from skin or eyes immediately.

AgriChem bv,

Industrial estate weststad

Koopvaardijweg 9 - 4906 CV Oosterhout (The Netherlands)

Postbus 295 - 4900 AG Oosterhout (The Netherlands)

Telephone: 00 31 162 431 931 Telefax: 00 31 162 456 797

E-mail: info@agrichem.com Webpage: www.agrichem.com



Vivendi[®] 200

MAPP 15017

WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Storage and disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank, and dispose of safely.

DO NOT RE-USE CONTAINER FOR ANY OTHER PURPOSE.

DIRECTIONS FOR USE

'IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.'

VIVENDI 200 is a post-emergence foliar acting herbicide. Although uptake can occur through roots and shoots VIVENDI 200 is taken up best by actively growing weed leaves. It is fully translocated throughout the plant and affects all parts of susceptible weeds. In mayweeds and thistles activity is greatest at the growing points, where death occurs first.

RESTRICTIONS

On the following food and fodder crops the statutory interval between the last application and harvest, consumption or conservation is the following:

Established grassland	- 7 days
Sugar beet, Fodder Beet, Mangel, Red beet	- 6 weeks
Swede, Turnip	- 6 weeks
Wheat, Barley & Oats	- Before GS 33 (GS 32 inclusive)
Oilseed rape	- apply before flower buds are visible from above crop canopy

Straw from crops treated with VIVENDI 200 should be baled and taken off the land. It can be burnt where this is permitted e.g. linseed. Do not use straw from crops treated with VIVENDI 200 for use in compost, manure or other use that may enable such straw to be used for any glasshouse or protected crop. Do not use manure from animals fed on crops treated with VIVENDI 200 for composting.

Do not apply VIVENDI 200 to crops undersown with clovers, sainfoin, vetches or other legumes.

Control of creeping thistle and other perennial weeds may be impaired by cultivations either before or within a period of two weeks after application of VIVENDI 200.

Do not apply to wet crops or when rain is expected within 6 hours of spraying.

Do not spray when crops are under stress from cold, drought, pest damage, nutrient deficiency etc.

Contract agents should be consulted before using on crops grown for seed.

Consult processors before using on crops for processing.

AgriChem bv,
Industrial estate weststad
Koopvaardijweg 9 - 4906 CV Oosterhout (The Netherlands)
Postbus 295 - 4900 AG Oosterhout (The Netherlands)
Telephone: 00 31 162 431 931 Telefax: 00 31 162 456 797
E-mail: info@agrichem.com Webpage: www.agrichem.com



Vivendi[®] 200

MAPP 15017

WEED CONTROL

VIVENDI 200 is active on a range of annual weeds, such as fat hen, but it should particularly be used for the control of thistles at a later stage.

Annual Weed Control

Unless otherwise specified, an application of 0.5 litres/ha VIVENDI 200 can be made to control annual weeds.

Weed	Susceptibility	Weed Growth Stage
Black-bindweed	MS	1 leaf
Corn marigold	S	2 leaf
Groundsel	S	6 leaf
Pale persicaria	MR	1 leaf
Pineappleweed	S	4 leaf
Redshank	MR	1 leaf
Mayweed spp	S	4 leaf
Smooth sow-thistle	S	2 leaf

S = susceptible; complete or near complete kill.

MS = moderately susceptible; good kill under favourable conditions.

MR = moderately resistant; poor kill but useful suppression.

Those weeds listed as susceptible may be controlled at a larger growth stage by the higher rate of use for perennial weed control.

Perennial Weed Control

Apply at 0.5 litre/ha. For the control of thistles a second application can be made at 1 litre/ha, which is usually 3-4 weeks after the first application. When thistles are present at the time of the application the dose of the first application can be increased to 1 litre/ha. However, please refer to the section 'CROP SPECIFIC INFORMATION' for advice on the different crops.

Weed	Susceptibility
Creeping thistle	S
Dandelion	S
Perennial sow-thistle	MS
Spear thistle	S

S = susceptible; complete or near complete kill.

MS = moderately susceptible; good kill under favourable conditions.

MR = moderately resistant; poor kill but useful suppression.

Weeds are best controlled when they are growing actively. The optimum timing of sprays targeted against thistles is at the 'rosette stage'.

Under cold, waterlogged or very dry conditions results will be reduced due to poorer plant growth in response to these circumstances.

Resistant weeds

Many weeds are resistant to VIVENDI 200. These include all Crucifer and Brassica weeds. They can however be controlled in tank mixture with other herbicides, provided the partner chemical is safe to the crop

AgriChem bv,
Industrial estate weststad
Koopvaardijweg 9 - 4906 CV Oosterhout (The Netherlands)
Postbus 295 - 4900 AG Oosterhout (The Netherlands)
Telephone: 00 31 162 431 931 Telefax: 00 31 162 456 797
E-mail: info@agrlichem.com Webpage: www.agrlichem.com



Vivendi[®] 200

MAPP 15017

being treated and is used in accordance with the partner label. Guidance is given on tank mixtures under the separate crop recommendations.

CROP SPECIFIC INFORMATION

1. CEREALS AND OILSEED RAPE

Winter and spring wheat, winter and spring barley and oats

For the control of annual weeds apply 0.35 litre/ha in the spring from second leaf stage (GS 12), to before the third node is detectable (up to and including GS 32).

VIVENDI 200 gives best results when applied with other herbicides in tank mixture provided the timings and conditions of use of the partner product coincide with those of VIVENDI 200.

Please refer to the section 'Compatibility' for further information about the possible mixing partners of VIVENDI 200.

Oilseed Rape (Winter or spring sown crops)

Apply 0.5 litre/ha in the spring for the control of annual weeds or 1.0 litre/ha in the spring for the control of creeping thistle, which is usually 3-4 weeks after the first application. Applications may be made from the two fully expanded leaf stage of the crop to before flower buds are visible at top of crop.

Please refer to the section 'Compatibility' for further information about the possible mixing partners of VIVENDI 200.

2. BEET CROPS AND FODDER CROPS

Sugar Beet, Fodder Beet, Red Beet and Mangel

VIVENDI 200 can be used once in repeat low dose herbicide programming on sugar beet, from the fully expanded cotyledon stage of the crop, in 80-100 litres/ha of water. Apply 0.5 litres/ha in tank mixture.

Where thistles are present an (additional) application of VIVENDI 200 alone at 1.0 litre/ha in 200-300 litres/ha water, may be used which is usually 3-4 weeks after the first application.

Please refer to the section 'Compatibility' for further information about the possible mixing partners of VIVENDI 200

Swede, Turnip

Apply 0.5 litre/ha for broad-leaved weed control once the crop has reached the two true leaf expanded stage.

Where thistles are a problem an (additional) application of 1.0 litre/ha can be made, which is usually 3-4 weeks after the first application.

Established Grassland

VIVENDI 200 can be used on permanent grassland and rotational grass that has been sown for at least one growing season. Livestock should not be left in fields to be sprayed. Application should be made when weeds are actively growing.

Apply 1.0 litre/ha. After grazing or cutting allow at least 2-3 weeks for regrowth of weeds before applying VIVENDI 200. An interval of 7 days must elapse after spraying VIVENDI 200 before livestock are re-introduced or the grass is cut for hay or silage.

AgriChem bv,
Industrial estate weststad
Koopvaardijweg 9 - 4906 CV Oosterhout (The Netherlands)
Postbus 295 - 4900 AG Oosterhout (The Netherlands)
Telephone: 00 31 162 431 931 Telefax: 00 31 162 456 797
E-mail: info@agrichem.com Webpage: www.agrichem.com



Vivendi[®] 200

MAPP 15017

3. HORTICULTURAL CROPS

Swede and Turnip

Apply 0.5 litre/ha for the control of annual weeds when seedling crops have at least two expanded true leaves and transplanted crops are fully established and have begun to make active growth.

Where thistles are a problem an (additional) application of 1.0 litre/ha can be made, which is usually 3-4 weeks after the first application.

Ornamental Plant Production

Applications must be only via directed sprays only to target weed foliage. Do not use between the end of August and beginning of March. Apply 1.0 litre/ha for the control of annual weeds or thistles. VIVENDI 200 can be sprayed at the optimum time for control of target weeds but it must only be applied to the weeds. Spray must be kept off ornamental plant foliage, buds, green wood or suckers, using a shielded sprayer if necessary. Do not apply in the root zone of the Compositae (e.g. Senecio) or Leguminosae (e.g. Laburnum, Genista) families.

FOLLOWING CROPS

Plant remains from crops treated with VIVENDI 200 may contain active clopyralid that could adversely affect succeeding crops especially peas, beans, lupins and other legumes carrots, parsnips, parsley and other umbellifers, potatoes, lettuce and other compositae, glasshouse and protected crops.

After crops treated with VIVENDI 200 (excluding cereals)

Do not plant susceptible crops in the same year as VIVENDI 200 has been applied or in the spring where VIVENDI 200 has been applied after July the previous year.

Chop and incorporate all treated plant remains as soon as possible after harvest to release residues into the soil where they are more quickly broken down. Ensure all plant remains from treated crops have fully decayed before planting susceptible crops.

After cereals treated with VIVENDI 200

Where straw from treated crops has been incorporated into the soil, only tolerant crops (e.g. cereals) should be drilled in the same year as treatment with VIVENDI 200.

MIXING AND SPRAYING

Mixing

Half fill the spray tank with clean water. If using in tank mixture add the mixture product first, in accordance with the mixture product label instructions, followed by the required quantity of VIVENDI 200. Add the remainder of the water and begin agitation. Continue agitation until spraying is completed.

Individual tank mix recommendations are given for separate crops. When VIVENDI 200 is to be used with other pesticides in tank mixture read the instructions of the partner product before mixing the spray. Ensure that the mixing instructions and conditions of use for the partner product are compatible with those of VIVENDI 200.

Spraying

Use a medium quality spray (BCPC definition).

Apply in a water volume of 200-300 litres/ha. Use the higher volume on larger weeds or where the crop is dense.

AgriChem bv,
Industrial estate weststad
Koopvaardijweg 9 - 4906 CV Oosterhout (The Netherlands)
Postbus 295 - 4900 AG Oosterhout (The Netherlands)
Telephone: 00 31 162 431 931 Telefax: 00 31 162 456 797
E-mail: info@agrichem.com Webpage: www.agrichem.com



Vivendi[®] 200

MAPP 15017

On very large weeds and in established grassland use 300-400 litres/ha.

After Spraying

VIVENDI 200 is active on susceptible plants and crops at very low doses, especially those of the Compositae and Leguminosae families. After spraying with VIVENDI 200 wash spray equipment thoroughly with water and a liquid detergent immediately after use. Spray out. Fill with clean water and leave overnight. Spray out again before storing or using another product. Dispose of the washings according to the official "Code of Practice for Farmers and Growers".

COMPATIBILITY

In cereals, oilseed rape and beet crops, VIVENDI 200 can be used in tank-mix with other approved pesticides.

VIVENDI 200 should be applied in recommended tank mix with other products only if any requirements or restrictions on the other product label do not conflict with the requirements or restrictions of VIVENDI 200.

Cereals

VIVENDI 200 can be tank mixed with the following approved formulations of:

MCPA salt	(MAPP 07400)
Mecoprop salt	(MAPP 14390, 13971, 14385)
Fluroxypyr	(MAPP 12018)

Oilseed Rape

VIVENDI 200 is compatible with the following approved formulations of:

Deltamethrin	(MAPP 07172)
Mancozeb	(MAPP 12564)

Beet

Conventional application:

VIVENDI 200 can be used in combination with approved formulations of:

Phenmedipham	(MAPP 15230, 13893, 14209, 14192)
Lenacil	(MAPP 09981)
Metamitron	(MAPP 11539, 14408, 12851, 13307)
Adjuvants Oils	(ADJ 0505, 0547)

Repeat low dose programme:

VIVENDI 200 can be used with approved formulations of:

Phenmedipham	(MAPP 15230, 13893, 14209, 14192)
Ethofumesate	(MAPP 12986, 12349)
Desmedipham + Ethofumesate + Phenmedipham	(MAPP 14034, 14262)
Metamitron	(MAPP 11539, 14408, 12851, 13307)
Adjuvant Oils	(ADJ 0505, 0547)

Use these tank mixtures in accordance with the crop timing and other conditions of use on the mixture partner's label.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and suitable for use. Since we cannot control the conditions of storage, handling or use or the weather conditions before or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality of fitness of

AgriChem bv,
Industrial estate weststad
Koopvaardijweg 9 - 4906 CV Oosterhout (The Netherlands)
Postbus 295 - 4900 AG Oosterhout (The Netherlands)
Telephone: 00 31 162 431 931 Telefax: 00 31 162 456 797
E-mail: info@agrichem.com Webpage: www.agrichem.com



Vivendi[®] 200

MAPP 15017

our goods are excluded. We cannot therefore accept any responsibility for any failure in performance, damage or injury whatsoever, arising from storage, handling, application or use.

PROTECT FROM FROST

AgriChem bv,
Industrial estate weststad
Koopvaardijweg 9 - 4906 CV Oosterhout (The Netherlands)
Postbus 295 - 4900 AG Oosterhout (The Netherlands)
Telephone: 00 31 162 431 931 Telefax: 00 31 162 456 797
E-mail: info@agrichem.com Webpage: www.agrichem.com

